thrombocytopenia-absent radius syndrome

Thrombocytopenia-absent radius (TAR) syndrome is characterized by the absence of a bone called the radius in each forearm and a shortage (deficiency) of blood cells involved in clotting (platelets). This platelet deficiency (thrombocytopenia) usually appears during infancy and becomes less severe over time; in some cases the platelet levels become normal.

Thrombocytopenia prevents normal blood clotting, resulting in easy bruising and frequent nosebleeds. Potentially life-threatening episodes of severe bleeding (hemorrhages) may occur in the brain and other organs, especially during the first year of life. Hemorrhages can damage the brain and lead to intellectual disability. Affected children who survive this period and do not have damaging hemorrhages in the brain usually have a normal life expectancy and normal intellectual development.

The severity of skeletal problems in TAR syndrome varies among affected individuals. The radius, which is the bone on the thumb side of the forearm, is almost always missing in both arms. The other bone in the forearm, which is called the ulna, is sometimes underdeveloped or absent in one or both arms. TAR syndrome is unusual among similar malformations in that affected individuals have thumbs, while people with other conditions involving an absent radius typically do not. However, there may be other abnormalities of the hands, such as webbed or fused fingers (syndactyly) or curved pinky fingers (fifth finger clinodactyly). Some people with TAR syndrome also have skeletal abnormalities affecting the upper arms, legs, or hip sockets.

Other features that can occur in TAR syndrome include malformations of the heart or kidneys. Some people with this disorder have unusual facial features including a small lower jaw (micrognathia), a prominent forehead, and low-set ears. About half of affected individuals have allergic reactions to cow's milk that may worsen the thrombocytopenia associated with this disorder.

Frequency

TAR syndrome is a rare disorder, affecting fewer than 1 in 100,000 newborns.

Genetic Changes

Mutations in the *RBM8A* gene cause TAR syndrome. The *RBM8A* gene provides instructions for making a protein called RNA-binding motif protein 8A. This protein is believed to be involved in several important cellular functions involving the production of other proteins.

Most people with TAR syndrome have a mutation in one copy of the *RBM8A* gene and a deletion of genetic material from chromosome 1 that includes the other copy of the *RBM8A* gene in each cell. A small number of affected individuals have mutations in both copies of the *RBM8A* gene in each cell and do not have a deletion on chromosome 1. *RBM8A* gene mutations that cause TAR syndrome reduce the amount of RNA-binding motif protein 8A in cells. The deletions involved in TAR syndrome eliminate at least 200,000 DNA building blocks (200 kilobases, or 200 kb) from the long (q) arm of chromosome 1 in a region called 1q21.1. The deletion eliminates one copy of the *RBM8A* gene in each cell and the RNA-binding motif protein 8A that would have been produced from it.

People with either an *RBM8A* gene mutation and a chromosome 1 deletion or with two gene mutations have a decreased amount of RNA-binding motif protein 8A. This reduction is thought to cause problems in the development of certain tissues, but it is unknown how it causes the specific signs and symptoms of TAR syndrome. No cases have been reported in which a deletion that includes the *RBM8A* gene occurs on both copies of chromosome 1; studies indicate that the complete loss of RNA-binding motif protein 8A is not compatible with life.

Researchers sometimes refer to the deletion in chromosome 1 associated with TAR syndrome as the 200-kb deletion to distinguish it from another chromosomal abnormality called a 1q21.1 microdeletion. People with a 1q21.1 microdeletion are missing a different, larger DNA segment in the chromosome 1q21.1 region near the area where the 200-kb deletion occurs. The chromosomal change related to 1q21.1 microdeletion is often called the recurrent distal 1.35-Mb deletion.

Inheritance Pattern

TAR syndrome is inherited in an autosomal recessive pattern, which means both copies of the gene in each cell are altered. In this disorder, either both copies of the *RBM8A* gene in each cell have mutations or, more commonly, one copy of the gene has a mutation and the other is lost as part of a deleted segment on chromosome 1. The affected individual usually inherits an *RBM8A* gene mutation from one parent. In about 75 percent of cases, the affected person inherits a copy of chromosome 1 with the 200-kb deletion from the other parent. In the remaining cases, the deletion occurs during the formation of reproductive cells (eggs and sperm) or in early fetal development. Although parents of an individual with TAR syndrome can carry an *RBM8A* gene mutation or a 200-kb deletion, they typically do not show signs and symptoms of the condition.

Other Names for This Condition

- chromosome 1q21.1 deletion syndrome, 200-KB
- radial aplasia-amegakaryocytic thrombocytopenia
- radial aplasia-thrombocytopenia syndrome

- TAR syndrome
- thrombocytopenia absent radii

Diagnosis & Management

Genetic Testing

 Genetic Testing Registry: Radial aplasia-thrombocytopenia syndrome https://www.ncbi.nlm.nih.gov/gtr/conditions/C0175703/

Other Diagnosis and Management Resources

- GeneReview: Thrombocytopenia Absent Radius Syndrome https://www.ncbi.nlm.nih.gov/books/NBK23758
- MedlinePlus Encyclopedia: Skeletal Limb Abnormalities https://medlineplus.gov/ency/article/003170.htm

General Information from MedlinePlus

- Diagnostic Tests https://medlineplus.gov/diagnostictests.html
- Drug Therapy https://medlineplus.gov/drugtherapy.html
- Genetic Counseling https://medlineplus.gov/geneticcounseling.html
- Palliative Care https://medlineplus.gov/palliativecare.html
- Surgery and Rehabilitation https://medlineplus.gov/surgeryandrehabilitation.html

Additional Information & Resources

MedlinePlus

- Encyclopedia: Skeletal Limb Abnormalities https://medlineplus.gov/ency/article/003170.htm
- Health Topic: Bleeding Disorders https://medlineplus.gov/bleedingdisorders.html

Genetic and Rare Diseases Information Center

 TAR syndrome https://rarediseases.info.nih.gov/diseases/5116/tar-syndrome

Additional NIH Resources

- National Cancer Institute (NCI): Inherited Bone Marrow Failure Syndromes https://dceg.cancer.gov/research/what-we-study/bone-marrow-failure-syndromes
- National Heart Lung and Blood Institute: What is Thrombocytopenia? https://www.nhlbi.nih.gov/health/health-topics/topics/thcp/
- National Human Genome Research Institute: Chromosome Abnormalities https://www.genome.gov/11508982/

Educational Resources

- American Society for Surgery of the Hand: Congenital Hand Differences http://www.assh.org/handcare/hand-arm-conditions/Congenital-Differences
- Boston Children's Hospital: Congenital Limb Defects http://www.childrenshospital.org/conditions-and-treatments/conditions/congenital-limb-defects
- Disease InfoSearch: TAR syndrome http://www.diseaseinfosearch.org/TAR+syndrome/6998
- MalaCards: thrombocytopenia-absent radius syndrome http://www.malacards.org/card/thrombocytopenia_absent_radius_syndrome
- Orphanet: Thrombocytopenia-absent radius syndrome http://www.orpha.net/consor/cgi-bin/OC_Exp.php?Lng=EN&Expert=3320

Patient Support and Advocacy Resources

- Birth Defect Research for Children http://www.birthdefects.org/thrombocytopenia-absent-radius-syndrome/
- Chromosome Disorder Outreach http://chromodisorder.org/
- Contact a Family http://www.cafamily.org.uk/medical-information/conditions/t/tar-syndrome/
- National Organization for Rare Disorders (NORD)
 https://rarediseases.org/rare-diseases/thrombocytopenia-absent-radius-syndrome/
- Resource List from the University of Kansas Medical Center http://www.kumc.edu/gec/support/tarsynd.html

GeneReviews

 Thrombocytopenia Absent Radius Syndrome https://www.ncbi.nlm.nih.gov/books/NBK23758

Scientific Articles on PubMed

PubMed

https://www.ncbi.nlm.nih.gov/pubmed?term=%28thrombocytopenia-absent+rad ius+syndrome%5BTIAB%5D%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+3600+days%22%5Bdp%5D

OMIM

 THROMBOCYTOPENIA-ABSENT RADIUS SYNDROME http://omim.org/entry/274000

Sources for This Summary

• Albers CA, Paul DS, Schulze H, Freson K, Stephens JC, Smethurst PA, Jolley JD, Cvejic A, Kostadima M, Bertone P, Breuning MH, Debili N, Deloukas P, Favier R, Fiedler J, Hobbs CM, Huang N, Hurles ME, Kiddle G, Krapels I, Nurden P, Ruivenkamp CA, Sambrook JG, Smith K, Stemple DL, Strauss G, Thys C, van Geet C, Newbury-Ecob R, Ouwehand WH, Ghevaert C. Compound inheritance of a low-frequency regulatory SNP and a rare null mutation in exon-junction complex subunit RBM8A causes TAR syndrome. Nat Genet. 2012 Feb 26;44(4):435-9, S1-2. doi: 10.1038/ng.1083.

Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/22366785

Free article on PubMed Central: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3428915/

- Bonsi L, Marchionni C, Alviano F, Lanzoni G, Franchina M, Costa R, Grossi A, Bagnara GP.
 Thrombocytopenia with absent radii (TAR) syndrome: from hemopoietic progenitor to mesenchymal stromal cell disease? Exp Hematol. 2009 Jan;37(1):1-7. doi: 10.1016/j.exphem.2008.09.004. Epub 2008 Nov 22. Review.
 - Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/19028006
- Greenhalgh KL, Howell RT, Bottani A, Ancliff PJ, Brunner HG, Verschuuren-Bemelmans CC, Vernon E, Brown KW, Newbury-Ecob RA. Thrombocytopenia-absent radius syndrome: a clinical genetic study. J Med Genet. 2002 Dec;39(12):876-81.
 Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/12471199
 Free article on PubMed Central: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1757221/
- Houeijeh A, Andrieux J, Saugier-Veber P, David A, Goldenberg A, Bonneau D, Fouassier M, Journel H, Martinovic J, Escande F, Devisme L, Bisiaux S, Chaffiotte C, Baux M, Kerckaert JP, Holder-Espinasse M, Manouvrier-Hanu S. Thrombocytopenia-absent radius (TAR) syndrome: a clinical genetic series of 14 further cases. impact of the associated 1q21.1 deletion on the genetic counselling. Eur J Med Genet. 2011 Sep-Oct;54(5):e471-7. doi: 10.1016/j.ejmg.2011.05.001. Epub 2011 May 13.
 - Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/21635976
- Klopocki E, Schulze H, Strauss G, Ott CE, Hall J, Trotier F, Fleischhauer S, Greenhalgh L, Newbury-Ecob RA, Neumann LM, Habenicht R, König R, Seemanova E, Megarbane A, Ropers HH, Ullmann R, Horn D, Mundlos S. Complex inheritance pattern resembling autosomal recessive inheritance involving a microdeletion in thrombocytopenia-absent radius syndrome. Am J Hum Genet. 2007 Feb;80(2):232-40. Epub 2006 Dec 21.

Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/17236129
Free article on PubMed Central: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1785342/

- Mokha J, Serrano M. Thrombocytopenia associated with cow's milk protein allergy: a case report.
 Clin Pediatr (Phila). 2013 Oct;52(10):985-7. doi: 10.1177/0009922812456593.
 Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/22914852
- Toriello HV. Thrombocytopenia-absent radius syndrome. Semin Thromb Hemost. 2011 Sep;37(6): 707-12. doi: 10.1055/s-0031-1291381. Epub 2011 Nov 18. Review.
 Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/22102274
- de Ybarrondo L, Barratt MS. Thrombocytopenia absent radius syndrome. Pediatr Rev. 2011 Sep; 32(9):399-400; discussion 400. doi: 10.1542/pir.32-9-399. Review.
 Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/21885665

Reprinted from Genetics Home Reference:

https://ghr.nlm.nih.gov/condition/thrombocytopenia-absent-radius-syndrome

Reviewed: February 2017 Published: March 21, 2017

Lister Hill National Center for Biomedical Communications U.S. National Library of Medicine National Institutes of Health Department of Health & Human Services